

Vortex solutions of parity and time reversal invariant Maxwell-Dirac-Chern-Simons gauge theory*

Junsoo Shin[†]

Department of Physics,

Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0435.

Abstract

We construct parity and time reversal invariant Maxwell-Chern-Simons gauge theory coupled to fermions with adding the parity partner to the matter and the gauge fields, which can give nontopological vortex solutions depending on the sign of the Chern-Simons coupling constant.

PACS numbers:11.15.-q, 11.15.Kc, 11.30.Er

Typeset using REVTeX

*Revised version (hep-th/9702124)

[†]Previous address: Department of Physics, Univ. of South Carolina, Columbia

Abelian gauge theories in (2+1)-dimensions with Chern-Simons terms have been found to admit interesting classical soliton solutions [1-5]. Especially, the fermionic field theories coupled to the Chern-Simons gauge field without Maxwell term admit vortex solutions [2]. Recently, it has been found that fermionic field theories coupled to the Maxwell-Chern-Simons gauge field also admit vortex solutions [3]. The Chern-Simons term has generally been believed to violate parity P and time reversal T [6]. However, it has been known that one can restore the parity and time reversal in Chern-Simons gauge theory by adding the parity partners to the matter and the gauge fields [7]. It is purpose of this work to show that such parity P and time reversal T invariant Maxwell-Dirac-Chern-Simons gauge theories can give the nontopological vortex solutions with binding energies.

We consider the Lagrangian

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}F_+^{\mu\nu}F_{+,\mu\nu} - \frac{1}{4}F_-^{\mu\nu}F_{-,\mu\nu} + \frac{\kappa}{4}\epsilon^{\mu\nu\rho}A_{+,\mu}F_{+,\nu\rho} - \frac{\kappa}{4}\epsilon^{\mu\nu\rho}A_{-,\mu}F_{-,\nu\rho} \\ & + i\bar{\psi}\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi + eA_{+,\mu}(J_+^\mu + lG_+^\mu) + eA_{-,\mu}(J_-^\mu - lG_-^\mu) \end{aligned} \quad (1)$$

where

$$F_\pm^{\mu\nu} = \partial^\mu A_\pm^\nu - \partial^\nu A_\pm^\mu, \quad J_\pm^\mu = \bar{\psi}\gamma^\mu P_\pm\psi, \quad G_\pm^\mu = \epsilon^{\mu\nu\rho}\partial_\nu J_{\pm,\rho}$$

and κ and l are coupling constants. The projectors $P_\pm$ onto the spin up and spin down fields are

$$P_\pm = \frac{1}{2}(1 \pm \rho_3),$$

where ρ_3 is a Dirac matrix defined by $\rho_i \equiv \sigma_i \otimes I$.

As in Ref. [3] and [8], we also introduce the topological current $G_{\pm,\mu}$, associated with the electromagnetic current $J_{\pm,\rho}$, which describes the induced charge and current density. The topological current takes the form,

$$G_\pm^i = \epsilon^{ij0}\partial_j J_{\pm,0} \quad (2)$$

which is related to the induced current by

$$G_{ind,\pm}^i = \pm l\epsilon^{ij0}\partial_j J_{\pm,0} \quad (3)$$

in the system [3] [8] . This induced current comes from the magnetic dipole moment density,

$$\vec{m}_{\pm} = \frac{\mu_{\pm}}{Q_{\pm}} J_{\pm,0} \hat{z} = \pm l J_{\pm,0} \hat{z} \quad (4)$$

through the relation $\vec{G}_{ind,\pm} = \nabla \times \vec{m}_{\pm}$ [3] [8] . Here, total charges are given by $Q_{\pm} = \int J_{\pm,0} d^2r$ and the magnetic dipole moments are $\mu_{\pm} \hat{z}$.

We choose the γ - matrices as

$$\gamma^0 = \begin{pmatrix} \sigma_3 & 0 \\ 0 & -\sigma_3 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} i\sigma_2 & 0 \\ 0 & i\sigma_2 \end{pmatrix}, \quad \gamma^2 = \begin{pmatrix} -i\sigma_1 & 0 \\ 0 & -i\sigma_1 \end{pmatrix}. \quad (5)$$

We write 4-component spinor field ψ as $\psi = \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix}$ where ψ_+ and ψ_- are two-component spinor functions.

The charge conjugation, parity and time-reversal transformation are defined as follows [9]:

1) Charge conjugation

$$\begin{aligned} A_{\pm,\mu}(x) &\rightarrow -A_{\pm,\mu}(x), \\ \psi_{\pm}(x) &\rightarrow \sigma_1 \psi_{\pm}^{\dagger}(x). \end{aligned} \quad (6)$$

2) Parity

$$\begin{aligned} A_{\pm,0}(t, x_1, x_2) &\rightarrow A_{\mp,0}(t, -x_1, x_2), \\ A_{\pm,1}(t, x_1, x_2) &\rightarrow \mp A_{\mp,1}(t, -x_1, x_2), \\ A_{\pm,2}(t, x_1, x_2) &\rightarrow A_{\mp,2}(t, -x_1, x_2), \\ \psi_{\pm}(t, x_1, x_2) &\rightarrow \sigma_1 \psi_{\mp}(t, -x_1, x_2). \end{aligned} \quad (7)$$

3) Time-reversal

$$\begin{aligned} A_{\pm,0}(t, x_1, x_2) &\rightarrow A_{\mp,0}(-t, x_1, x_2), \\ A_{\pm,i}(t, x_1, x_2) &\rightarrow -A_{\mp,i}(-t, x_1, x_2) \quad (i = 1, 2), \\ \psi_{\pm}(t, x_1, x_2) &\rightarrow \pm \sigma_1 \psi_{\mp}^*(-t, x_1, x_2). \end{aligned} \quad (8)$$

With this definition the Lagrangian density (1) is invariant under the parity and time-reversal transformation as well as PCT.

The equations of motion are

$$\partial_\nu F_\pm^{\nu\mu} \pm \frac{\kappa}{2} \epsilon^{\mu\nu\rho} F_{\pm,\nu\rho} = -e(J_\pm^\mu \pm lG_\pm^\mu), \quad (9)$$

and

$$\begin{aligned} & \gamma^\mu (i\partial_\mu + eP_+ A_{+,\mu} + eP_- A_{-,\mu})\psi - m\psi \\ & -el\epsilon^{\mu\nu\rho}(\partial_\nu A_{+,\mu})\gamma_\rho P_+ \psi + el\epsilon^{\mu\nu\rho}(\partial_\nu A_{-,\mu})\gamma_\rho P_- \psi = 0. \end{aligned} \quad (10)$$

We choose the gauge $A_{\pm,0} = 0$ and consider the gauge field $A_{\pm,i}$ to be static. By taking the fermion fields ψ_+ and ψ_- in component form,

$$\psi_+ = \begin{pmatrix} \eta_+ \\ \xi_+ \end{pmatrix} e^{-iE_+ t}, \quad \psi_- = \begin{pmatrix} \eta_- \\ \xi_- \end{pmatrix} e^{-iE_- t},$$

the equations of motion (10) can be written as

$$\begin{aligned} [E_+ - m - el\epsilon^{ij0}\partial_j A_{+,i}]\eta_+ &= -D_{++}\xi_+ + it[(i\partial_1 + \partial_2)E_+]\xi_+, \\ [-E_+ - m + el\epsilon^{ij0}\partial_j A_{+,i}]\xi_+ &= -D_{+-}\eta_+ + it[(-i\partial_1 + \partial_2)E_+]\eta_+ \end{aligned} \quad (11)$$

and

$$\begin{aligned} [-E_- - m - el\epsilon^{ij0}\partial_j A_{-,i}]\eta_- &= -D_{-+}\xi_- + it[(i\partial_1 + \partial_2)E_-]\xi_-, \\ [E_- - m + el\epsilon^{ij0}\partial_j A_{-,i}]\xi_- &= -D_{--}\eta_- + it[(-i\partial_1 + \partial_2)E_-]\eta_- \end{aligned} \quad (12)$$

where $D_{+\pm} = \pm iD_1^+ + D_2^+$, $D_{-\pm} = \pm iD_1^- + D_2^-$ and $D_i^\pm = \partial_i - ieA_{\pm,i}$.

If we let

$$\psi_+ = \eta_+ \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{-iE_+ t}, \quad \psi_- = \xi_- \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^{-iE_- t}, \quad (13)$$

the equations of motion (11) and (12) become

$$\begin{aligned}
(E_+ - m - el\epsilon^{ij0}\partial_j A_{+,i})\eta_+ &= 0, \\
D_{+-}\eta_+ - it[(-i\partial_1 + \partial_2)E_+]\eta_+ &= 0
\end{aligned} \tag{14}$$

and

$$\begin{aligned}
(E_- - m + el\epsilon^{ij0}\partial_j A_{-,i})\xi_- &= 0, \\
D_{-+}\xi_- - it[(i\partial_1 + \partial_2)E_-]\xi_- &= 0.
\end{aligned} \tag{15}$$

As shown in Ref. [3], from the form of the solutions (13), we find that the electromagnetic current and the induced charge vanish;

$$J_{\pm}^i = \bar{\psi}\gamma^i P_{\pm}\psi = 0, \quad G_{\pm}^0 = \epsilon^{0ij}\partial_i J_{\pm,j} = 0. \tag{16}$$

By using Eqs.(9) and (16), the magnetic fields reduce to

$$B_{\pm} = -F_{\pm,12} = \pm \frac{e}{\kappa} \rho_{\pm} \tag{17}$$

where $\rho_{\pm} \equiv \psi_{\pm}^{\dagger}\psi_{\pm}$. We note that Eqs.(9) can be written as

$$\partial_j F_{\pm}^{ji} = \mp el\epsilon^{ij0}\partial_j J_{\pm,0}. \tag{18}$$

From Eqs.(17) and (18), We can find the constant l to be $l = -\frac{1}{\kappa}$.

To find the general solutions for the self-dual equation (14) and (15), we note when η_+ and ξ_- are decomposed into its phase and amplitude,

$$\eta_+ = \sqrt{\rho_+}e^{i\omega_+}, \quad \xi_- = \sqrt{\rho_-}e^{i\omega_-}, \tag{19}$$

the Eqs.(14) and (15) can be written as

$$e\nabla \times \vec{A}_{\pm} = \nabla \times (\nabla\omega_{\pm}) - \nabla \times (\nabla E_{\pm})t \mp \frac{1}{2}\nabla^2 \ln \rho_{\pm}. \tag{20}$$

From Eqs.(17) and (20) we obtain the equations for the charge density $\rho_{\pm}$:

$$\nabla^2 \ln \rho_{\pm} = -\frac{2e^2}{\kappa} \rho_{\pm}. \tag{21}$$

Eqs.(21) are the Liouville equations which are completely integrable. When we take the solutions (13), $\kappa > 0$ is required in order to have nonsingular positive solutions for the density ρ_+ and ρ_- . The most general circularly symmetric nonsingular solutions to the Liouville equations involve two positive constants $r_\pm$ and $\mathcal{N}_\pm$ [4] :

$$\rho_\pm = \frac{4\kappa\mathcal{N}_\pm^2}{e^2r^2} \left[\left(\frac{r_\pm}{r} \right)^{\mathcal{N}_\pm} + \left(\frac{r}{r_\pm} \right)^{\mathcal{N}_\pm} \right]^{-2} \quad (22)$$

where $r_\pm$ are scale parameters. In case of solutons (13), the magnetic dipole moments are given by $\mu_\pm = \pm lQ_\pm = \mp \frac{Q_\pm}{\kappa} (\kappa > 0)$.

If we let

$$\psi_+ = \eta_+ \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{-iE_+t}, \quad \psi_- = \eta_- \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{-iE_-t}, \quad (23)$$

the equations of motion (11) and (12) become

$$\begin{aligned} (E_+ - m - el\epsilon^{ij0}\partial_j A_{+,i})\eta_+ &= 0, \\ D_{+-}\eta_+ - it[(-i\partial_1 + \partial_2)E_+]\eta_+ &= 0 \end{aligned} \quad (24)$$

and

$$\begin{aligned} (E_- + m + el\epsilon^{ij0}\partial_j A_{-,i})\eta_- &= 0, \\ D_{--}\eta_- - it[(-i\partial_1 + \partial_2)E_-]\eta_- &= 0. \end{aligned} \quad (25)$$

If $\eta_\pm$ are decomposed into its phase and amplitude,

$$\eta_+ = \sqrt{\rho_+} e^{i\theta_+}, \quad \eta_- = \sqrt{\rho_-} e^{i\theta_-}, \quad (26)$$

the amplitudes $\rho_\pm$ satisfy the Liouville equations

$$\nabla^2 \ln \rho_\pm = \mp \frac{2e^2}{\kappa} \rho_\pm \quad (27)$$

where the $\mp$ signs are for the solutions of the forms $\psi_\pm$, respectively. Eqs.(27) imply that there is no nonsingular positive solution to satisfy the Liouville equations with respect to $\rho_\pm$ simultaneously.

If we let

$$\psi_+ = \xi_+ \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^{-iE_+t}, \quad \psi_- = \eta_- \begin{pmatrix} 1 \\ 0 \end{pmatrix} e^{-iE_-t}, \quad (28)$$

the equations of motion (11) and (12) become

$$\begin{aligned} (E_+ + m - el\epsilon^{ij0}\partial_j A_{+,i})\xi_+ &= 0, \\ D_{++}\xi_+ - it[(i\partial_1 + \partial_2)E_+]\xi_+ &= 0 \end{aligned} \quad (29)$$

and

$$\begin{aligned} (E_- + m + el\epsilon^{ij0}\partial_j A_{-,i})\eta_- &= 0, \\ D_{--}\eta_- - it[(-i\partial_1 + \partial_2)E_-]\eta_- &= 0. \end{aligned} \quad (30)$$

From the form of the solutions (28), the electromagnetic current $J_\pm^i$, and the induced charge $G_\pm^0$ also vanish. Therefore, the magnetic fields reduce to

$$B_\pm = -F_{\pm,12} = \pm \frac{e}{\kappa} \rho_\pm \quad (31)$$

and the constant l to be $l = -\frac{1}{\kappa}$.

In this case, we obtain the Liouville equations for the charge density $\rho_\pm$:

$$\nabla^2 \ln \rho_\pm = \frac{2e^2}{\kappa} \rho_\pm. \quad (32)$$

As for Eqs.(32), $\kappa < 0$ is required in order to have nonsingular positive charge density $\rho_\pm$. In this case, the most general circularly symmetric nonsingular solutions to the Liouville equations involve two positive constants $r_\pm$ and $\mathcal{N}_\pm$ [4] :

$$\rho_\pm = -\frac{4\kappa\mathcal{N}_\pm^2}{e^2 r^2} \left[\left(\frac{r_\pm}{r} \right)^{\mathcal{N}_\pm} + \left(\frac{r}{r_\pm} \right)^{\mathcal{N}_\pm} \right]^{-2} \quad (33)$$

In case of solutions (28), the magnetic dipole moments are given by $\mu_\pm = \mp \frac{Q_\pm}{\kappa} (\kappa < 0)$.

If we let

$$\psi_+ = \xi_+ \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^{-iE_+t}, \quad \psi_- = \xi_- \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^{-iE_-t}, \quad (34)$$

the equations of motion (11) and (12) become

$$\begin{aligned}(E_+ + m - el\epsilon^{ij0}\partial_j A_{+,i})\xi_+ &= 0, \\ D_{++}\xi_+ - it[(i\partial_1 + \partial_2)E_+]\xi_+ &= 0\end{aligned}\tag{35}$$

and

$$\begin{aligned}(E_- - m + el\epsilon^{ij0}\partial_j A_{-,i})\xi_- &= 0, \\ D_{--}\eta_- - it[(i\partial_1 + \partial_2)E_-]\xi_- &= 0.\end{aligned}\tag{36}$$

In this case, we obtain the Liouville equations for the charge density $\rho_{\pm}$:

$$\nabla^2 \ln \rho_{\pm} = \pm \frac{2e^2}{\kappa} \rho_{\pm}.\tag{37}$$

As for Eqs.(37), there is no nonsingular positive solution to satisfy the Liouville equations.

In cases of solutions (13) and (28), the Hamiltonian density $\mathcal{H}$ reduces to

$$\mathcal{H} = \frac{1}{2}F_{+,12}^2 + \frac{1}{2}F_{-,12}^2 \pm m\rho_+ \pm m\rho_- + \frac{e^2 l}{\kappa}\rho_+^2 + \frac{e^2 l}{\kappa}\rho_-^2,\tag{38}$$

respectively. The total energy of the system can be written as

$$\begin{aligned}E &= \int d^2r \mathcal{H} = \int d^2r [\pm m(\rho_+ + \rho_-) - \frac{e^2}{2\kappa^2}(\rho_+^2 + \rho_-^2)] \\ &= \pm m \frac{2\kappa}{e^2}(\mathcal{N}_+ + \mathcal{N}_-) \\ &\quad - \frac{e^2}{\kappa^2} \left[\left(\frac{4\pi\kappa^2 \mathcal{N}_+^3 \Gamma(2 - \frac{1}{\mathcal{N}_+}) \Gamma(2 + \frac{1}{\mathcal{N}_+})}{3e^4 r_+^2} \right) + \left(\frac{4\pi\kappa^2 \mathcal{N}_-^3 \Gamma(2 - \frac{1}{\mathcal{N}_-}) \Gamma(2 + \frac{1}{\mathcal{N}_-})}{3e^4 r_-^2} \right) \right].\end{aligned}\tag{39}$$

As in Ref. [3], $(\mathcal{N}_+, \mathcal{N}_-)$ -soliton solutions of this model (1) also have binding energies. Thus the stability of the nontopological $(\mathcal{N}_+, \mathcal{N}_-)$ -solitons in this model is guaranteed by the binding energies and the charge conservation laws of the system [3].

From the Eqs.(14) and (15), we find

$$E_+ = m - \frac{e^2}{\kappa^2}\rho_+, \quad E_- = m - \frac{e^2}{\kappa^2}\rho_-.\tag{40}$$

From the Eqs.(29) and (30), we also find

$$E_+ = -m - \frac{e^2}{\kappa^2}\rho_+, \quad E_- = -m - \frac{e^2}{\kappa^2}\rho_-. \quad (41)$$

Therefore, the solutions (13) or (28) is acceptable because the parity transformation does not change any internal properties of the particle like the energy except for the binding energy [5]. However, from the Eqs.(24) and (25), we find

$$E_+ = m - \frac{e^2}{\kappa^2}\rho_+, \quad E_- = -m - \frac{e^2}{\kappa^2}\rho_-. \quad (42)$$

From Eqs.(35) and (36), we also find

$$E_+ = -m - \frac{e^2}{\kappa^2}\rho_+, \quad E_- = m - \frac{e^2}{\kappa^2}\rho_-. \quad (43)$$

In these cases, we find that the definition of Eqs.(7) changes the internal property under the parity transformation. In other words, the parity transformation implies that each solution transforms to the solution with the different energy. This is unacceptable because the parity transformation should not change any internal properties of the particles [5]. Therefore, we can also ascertain that the forms (23) and (34) are not the solutions of this model (1).

ACKNOWLEDGMENTS

The author has benefited from conversations with J.M. Knight, L.N. Chang and J.H. Yee and would like to thank Feng-Li Lin for helpful discussion.

REFERENCES

- [1] S. Deser, R. Jackiw, and S. Templeton, Ann. Phys. (N.Y.) **140**, 372 (1982); J. Hong, Y. Kim, and P.Y. Pac, Phys. Rev. Lett. **64**, 2230 (1990); R. Jackiw and E.J. Weinberg, ibid. **64**, 2234 (1990); R. Jackiw, K. Lee and Erick J. Weinberg, Phys. Rev. **D42**, 3488 (1990); C. Lee, K. Lee and H. Min, Phys. Lett. **B252**, 79 (1990).
- [2] Y.M. Cho, J.W. Kim, and D.H. Park, Phys. Rev. **D45**, 3802 (1992); C. Duval, P.A. Horvathy, and L. Palla, ibid. **52**, 4700 (1995).
- [3] S. Hyun, J. Shin, J.H. Yee and H. Lee, Phys. Rev. **D55**, 3900 (1997).
- [4] R. Jackiw and S.Y. Pi, Phys. Rev. Lett. **64** 2969 (1990); R. Jackiw and S.Y. Pi, Phys. Rev. **D42** 3500 (1990).
- [5] J. Shin and J.H. Yee, Phys. Rev. **D50**, 4223 (1994).
- [6] J. Fröhlich and A. Zee, Nucl. Phys. **B364**, 517 (1991); K.B. Lyons, J. Kwo, J.F. Dillion, G.P. Espinosa, M. McGlashan-Powell, A. Ramirez, and L. Schneemeyer, Phys. Rev. Lett. **64**, 2949 (1990); S. Spielman, K. Fesler, C.B. Eom, T.H. Geballe, M.M. Fejer, and A. Kapitulnik, Phys. Rev. Lett. **65**, 123 (1990); H.J. Weber, D. Weitbrecht, D. Brach, A.L. Shelankov, H. Keiter, W. Weber, Th. Wolf, J. Geerk, G. Linker, G. Roth, P.C. Splittgerber-Hünnekes, and G. Güntherodt, Solid State Commun. **76**, 511 (1990) .
- [7] J. Schonfeld, Nucl. Phys. **B185**, 157 (1981) ; G. Semenoff and N. Weiss, Phys. Lett. **B250**, 117 (1990) ; C.R. Hagen, Phys. Rev. Lett. **68**, 3821 (1992) .
- [8] D.G. Barci and L.E. Oxman, Phys. Rev. **D52**, 1169 (1995); M.E. Carrington and G. Kunstatter, ibid. **51**, 1903 (1995).
- [9] Y.M. Cho, SNUTP 93-01 Preprint (1993)